



38077

Hospital identification number 0 1 2 3 4 5 6 7 8 9

European Pressure Ulcer Prevalence Study Minimum Data Set



General Data

- Setting** University Hospital General Hospital Local Hospital
- Country** Belgium Italy Denmark Netherlands Finland Portugal France Spain Germany Sweden Greece United Kingdom Hungary Other
- Number of Beds** > 1000 500 - 1000 < 500

Patient Data

- Age** < 12 12 - 18 19 - 39 40 - 59 60 - 69 70 - 79 80 - 89 > 89
- Gender** Female Male
- Care Group** Neurology / Rehabilitation Intensive Chronic Care Acute Care / High Dependency
- Expected Length of Stay** < 6 days 6 days - 1 month > 1 month

Braden Scale

| | | |
|--|---|---|
| Sensory perception <input type="radio"/> Completely limited <input type="radio"/> Very limited <input type="radio"/> Slightly limited <input type="radio"/> No impairment | Moisture <input type="radio"/> Constantly moist <input type="radio"/> Very moist <input type="radio"/> Occasionally moist <input type="radio"/> Rarely moist | Activity <input type="radio"/> Bedfast <input type="radio"/> Chairfast <input type="radio"/> Walks occasionally <input type="radio"/> Walks frequently |
| Mobility <input type="radio"/> Completely immobile <input type="radio"/> Very limited <input type="radio"/> Slightly limited <input type="radio"/> No limitations | Nutrition <input type="radio"/> Very poor <input type="radio"/> Probably inadequate <input type="radio"/> Adequate <input type="radio"/> Excellent | Friction and Shear <input type="radio"/> Problem <input type="radio"/> Potential problem <input type="radio"/> No apparent problem |

Incontinence

- Incontinence**
 Not
 Occasional
 Usually/Urine
 Double

Skin observation


Most severe pressure ulcer

- None
 Non-blanchable erythema
 Blister / Abrasion
 Superficial ulcer
 Deep ulcer / Necrosis

Location(s) of most severe pressure ulcer(s)

- Sacrum
 Heel
 Hip
 Other

All existing pressure ulcers



Prevention

- Equipment**
- In bed** No special equipment Non-powered device Powered device
- In chair** No special equipment Non-powered device Powered device
- Repositioning**
- In bed** No planned/Irregularly Every 2 hours Every 3 hours Every 4 hours
- In chair** No planned/Irregularly Every 2 hours Every 3 hours Every 4 hours